

# 鳥取大学研究成果リポジトリ

## Tottori University research result repository

タイトル Title	Trichilemmal cyst with dermatofibroma-like features on the leg
著者 Author(s)	Ehara, Yuko; Yoshida, Yuichi; Yanagihara, Shigeto; Shiomi, Tatsushi; Yamamoto, Osamu
掲載誌・巻号・ページ Citation	JOURNAL OF DERMATOLOGY , 44 (4) : E71 - E72
刊行日 Issue Date	2017-04
資源タイプ Resource Type	その他 / Others
版区分 Resource Version	著者版 / Author
権利 Rights	(C) 2016 Japanese Dermatological Association.
DOI	<a href="https://doi.org/10.1111/1346-8138.13573">10.1111/1346-8138.13573</a>
URL	<a href="https://repository.lib.tottori-u.ac.jp/8010">https://repository.lib.tottori-u.ac.jp/8010</a>

**Title: Trichilemmal cyst with dermatofibroma-like features on the leg**

Short title: Trichilemmal cyst with dermatofibroma-like features

The total number of words 463 figures 1(1a, 1b, 1c, 1d, 1e, 1f and 1g) and tables(0) and references 5.

Yuko EHARA<sup>1</sup>, Yuichi YOSHIDA<sup>1</sup>, Shigeto YANAGIHARA<sup>1</sup>, Tatsushi SHIOMI<sup>2</sup>,  
Osamu YAMAMOTO<sup>1</sup>

<sup>1</sup>Division of Dermatology, Department of Medicine of Sensory and Motor Organs, and

<sup>2</sup>Organ Pathology, Faculty of Medicine, Tottori University, Yonago, Japan

Correspondence to: Yuko Ehara

Division of Dermatology, Department of Medicine of Sensory and Motor Organs,  
Faculty of Medicine, Tottori University, 86 Nishi-cho, Yonago-shi, Tottori 683-8503,  
Japan

Tel: +81-859-38-6597      Fax: +81-859-38-6599

Email: [yukoehara@med.tottori-u.ac.jp](mailto:yukoehara@med.tottori-u.ac.jp)

Funding sources: none, Conflict of interest: none

(Key words) trichilemmal cyst, pilar cyst, dermatofibroma, dermoscopy.

We report here a case of TC showing both dermoscopic and histopathologic features similar to dermatofibroma (DF).

A 71-year-old Japanese woman was referred to our department for evaluation of a 1-year history of nodule on her leg. Physical examination revealed a brownish nodule, 9 × 6 mm in diameter, located on the right lower leg (Fig. 1a). Dermoscopic examination showed a central yellowish-white patch with a peripheral pigment network (Fig. 1b).

Histopathological examination showed an intradermal cyst that contained keratinous materials. The cells in the peripheral layer of the wall exhibited a palisading arrangement without a granular layer (Fig. 1c and 1d). Elongation of rete ridges and hyperpigmentation in the epidermis were also observed (Fig. 1e), and there was an increase in dermal collagen bundles with inflammatory cells and multinucleated histiocytes around the cyst reminiscent of DF. Immunohistochemically, the inflammatory cells around the cyst were positive for CD10 (Fig. 1f) and CD68 (Fig. 1g).

Finally, a diagnosis of trichilemmal cyst with DF-like features was made.

TC mostly occurs on the scalp, but about 1.4% of TCs were seen on extremities.<sup>1</sup>

Histopathological findings of our case were typical for TC with DF-like features. To our knowledge, there has been no report of TC with DF-like features in both histopathologic and dermoscopic aspects.

Immunohistological staining with CD10 and/or CD68 is usually useful for diagnosis of DF.<sup>2, 3</sup> When TC undergoes partial or complete disintegration, an endogenous foreign-body reaction is induced. We considered that the DF-like features were derived from infiltration of inflammatory cells and fibroplasia.

With respect to dermoscopic examination, there has been only one case in which dermoscopic findings of TC was reported. Gencoglan et al.<sup>4</sup> reported that diffuse homogeneous blue pigmentation with a minimal reddish hue was seen in TC on the arm. They considered that these features were due to homogeneous dense keratin materials, so-called Tyndall effect.<sup>4</sup> However, dermoscopic findings may vary by anatomical location and depth of the tumor.

Dermoscopic findings of DF are well-known. The most common dermoscopic pattern is a central white patch with a delicate pigment network. The former corresponds to pronounced fibrosis within the papillary dermis, and the latter is a result of hyperpigmentation of basal cells in the epidermis.<sup>5</sup>

In our case, dermoscopic examination showed a pattern similar to that of DF. It is likely that the central white area was derived from keratinous materials and inflammatory cells around the cyst. The delicate pigment network at the periphery corresponded to hyperpigmentation in the basal layer. No reddish hue was seen in our case.

TCs are commonly observed on the scalp, but there have been few reports on dermoscopic observations of TC. In our case, the clinical features were not typical of TC. Dermoscopy may provide additional clues for a diagnosis in non-typical skin tumors.

**Disclosure.** *Financial support: none. Conflict of interest: none.*

## REFERENCES

1. McGauran MH, Binnington B. Keratinous cysts of the skin: identification and differentiation of pilar cysts from epidermal cysts. *Arch Dermatol* 1966; **94**: 499-508.
2. Clarke LE, Frauenhoffer E, Fox E, Neves R, Bruggeman RD, Helm KF. CD10-positive myofibrosarcomas: a pitfall in the differential diagnosis of atypical fibroxanthoma. *J Cutan Pathol* 2010; **7**: 737-43.
3. Pulford KA, Rigney EM, Micklem KJ, et al. KP1: a new monoclonal antibody that detects a monocyte/macrophage associated antigen in routinely processed tissue sections. *J Clin Pathol* 1989; **4**: 414-21.
4. Gencoglan G, Karaarslan IK, Akalin T, Ozdemir F. Trichilemmal cyst with homogeneous blue pigmentation on dermoscopy. *Australasian Journal of Dermatology* 2009; **50**: 301-302.
5. Puig S, Romero D, Zaballos P, Malvehy J. Dermoscopy of Dermatofibroma. *Arch Dermatol* 2005; **1**: 122.

## Figure Legends

**Figure 1.** (a) A brownish nodule on the right lower leg. (b) The lesion showed a central white patch with a peripheral delicate pigment network by dermoscopy. (c, d) The wall cells of the peripheral layer showed trichilemmal keratinization (haematoxylin & eosin (HE), original magnification  $\times 20$  and  $\times 100$ ). (e) Dermatofibroma-like features were seen (HE  $\times 20$  and  $\times 100$ ). (f, g) Immunostaining was positive for CD10 and CD68 ( $\times 100$ ).



